



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}
\therefore V &= \int_{ch/R}^h \left\{ \frac{R^2 x^2}{h^2} \cos^{-1} \left(\frac{ch}{Rx} \right) - \frac{c}{h} \sqrt{R^2 x^2 - h^2 c^2} \right\} dx, \\
&= \frac{1}{3} h \left\{ R^2 \cos^{-1} \left(\frac{c}{R} \right) - 2c \sqrt{R^2 - c^2} + \frac{c^3}{R} \log \left(\frac{R + \sqrt{R^2 - c^2}}{c} \right) \right\}, \\
&= 3 \left\{ 9 \cos^{-1} \left(\frac{1}{\sqrt{2}} \right) - 9 + \frac{9\sqrt{2}}{4} \log(\sqrt{2} + 1) \right\}, = 2.619 \text{ cubic inches.}
\end{aligned}$$

III. Solution by J. SCHEFFER, A. M., Hagerstown, Md.

I Solution. Designating, the radius of the base by r , the altitude by h , and choosing the center of the base for the origin of orthogonal coördinates, CO for the axis of z , the radius OB for the axis of x and a radius parallel to the section FDG for that of y , we find the equation of the cone to be

$$z = (h/r)(r - \sqrt{x^2 + y^2}),$$

and the volume V of $COFGD$

$$\begin{aligned}
&= \int_{-\sqrt{r^2-x^2}}^{-\sqrt{r^2+x^2}} \frac{h}{r} dy \int_0^x dx \int_{-\sqrt{r^2-x^2}}^{\sqrt{r^2-x^2}} (r - \sqrt{x^2 + y^2}) dy, \\
&= \frac{h}{r} \int_0^r \left(r \sqrt{r^2 - x^2} - x^2 \log \frac{r + \sqrt{r^2 - x^2}}{x} \right) dx, \\
&= \frac{2}{3} h x \sqrt{r^2 - x^2} + \frac{1}{3} r^2 h \sin^{-1}(x/r) - \frac{1}{3} \frac{h x^3}{r} \log \frac{r + \sqrt{r^2 - x^2}}{x}.
\end{aligned}$$

Substituting $x = (r/2)\sqrt{2}$, we have for the volume $COFGD$ the expression $\frac{r^2 h}{12} [4 + \pi - \sqrt{2} \cdot \log(\sqrt{2} + 1)]$, and for that of $B-FGD$ $\frac{r^2 h}{12} [\pi - 4 + \sqrt{2} \cdot \log(\sqrt{2} + 1)]$

II Solution. Let HK be a circle parallel to AB cutting the hyperbola FDK in the points L and M , and let the diameter HK cut the axis DE at Q . Put $OE = b$, $OF = r$, $CO = h$, $DQ = x$, $LQ = y$. We find from the geometry of the figure $y^2 = (2br/x)x + (r^2/h^2)x^2$ as the equation of the hyperbola FDK .

$$\therefore \text{Area of } FDK = 2 \int dx \sqrt{\frac{2br}{h}x + \frac{r^2}{h^2}x^2} \text{ between the limits } O,$$

and $DE = \frac{(r-b)h}{r}$. Integrating we find for this area the expression,

$$\frac{h}{b} \left(r \sqrt{r^2 - b^2} - b^2 \log \frac{r + \sqrt{r^2 - b^2}}{b} \right).$$

$$\therefore \text{Volume of } COFGD = \frac{h}{b} \int_0^b \left(r \sqrt{r^2 - b^2} - b^2 \log \frac{r + \sqrt{r^2 - b^2}}{b} \right) db$$

$$= \frac{2}{3}hb\sqrt{r^2 - b^2} + \frac{1}{3}r^2h\sin^{-1}\frac{b}{r} - \frac{1}{3}\frac{hb^3}{r}\log\frac{r+\sqrt{r^2-b^2}}{b},$$

$$\text{and volume of } BFGD = \frac{1}{3}r^2h\cos^{-1}\frac{b}{r} - \frac{2}{3}hb\sqrt{r^2 - b^2} + \frac{1}{3}\frac{hb^3}{r}\log\frac{r+\sqrt{r^2-b^2}}{b}.$$

III Solution. Let HK be a circle parallel to AB , and N its centre. Through N draw a diameter parallel to the hyperbolic section FGD . Put $CN=x$, $OE=b$, $BO=r$, $CO=h$, then the area of the circular segment lying between the diameter through N and the parallel chord LM

$$= \frac{r^2x^2}{h^2}\sin^{-1}\frac{bh}{rx} + \frac{br}{h}\sqrt{x^2 - \frac{b^2h^2}{r^2}}.$$

\therefore Volume of conical section $COFGD$

$$= \frac{r}{h} \left\{ \frac{r}{h} \int x^2 \sin^{-1} \frac{bh}{rx} + b \int dx \sqrt{x^2 - \frac{b^2h^2}{r^2}} \right\},$$

the integrals to be taken between $h-DE=bh/r$ and h . Thus we find for this volume the expression

$$\frac{2}{3}hb\sqrt{r^2 - b^2} + \frac{1}{3}r^2h\sin^{-1}\frac{b}{r} - \frac{1}{3}\frac{hb^3}{r}\log\frac{r+\sqrt{r^2-b^2}}{r};$$

and for the volume of the conical section $DBFG$,

$$\frac{1}{3}r^2h\cos^{-1}\frac{b}{r} - \frac{2}{3}hb\sqrt{r^2 - b^2} + \frac{1}{3}\frac{hb^3}{r}\log\frac{r+\sqrt{r^2-b^2}}{r}.$$

HISTORICAL NOTE. The famous astronomer KEPLER tried hard to find the volume of such conical sections as the above, but all his efforts proved futile.

Also solved by *GEORGE LILLEY, Ph. D., LL. D.*, and *CHARLES C. CROSS*. Dr. Lilley obtained a numerical result of 6.771 cubic inches, and Professor Cross obtained 2.256979 cubic inches.

PROBLEMS FOR SOLUTION.

ARITHMETIC.

87. Proposed by *E. W. MORRELL, A. M.*, Professor of Mathematics, Montpelier Seminary, Montpelier, Vt.

A and B set out from the same place, and in the same direction. A travels uniformly 18 miles per day, and after 9 days turns and goes back as far as B has traveled during those 9 days; he then turns again, and, pursuing his journey, overtakes B 22½ days after the time they first set out. It is required to find the rate at which B uniformly traveled. [From *Greenleaf's Arithmetic*.]

88. Proposed by *J. A. CALDERHEAD, M. Sc.*, Professor of Mathematics, Curry University, Pittsburg, Pa.

Find the principal of a note given March 19, 1891, bearing interest at 6%. Payments: September 1, 1892, \$243.50; January 19, 1893, \$6.90; April 13, 1894, \$19.10; September 19, 1894, \$110.90. Amount due February 22, 1897, \$229.10.